



HOW TO USE GREEN BULL ONE TREATMENT:

The information in these guidelines may be different depending on the country-specific climate and crops.

In the case of extreme climatic conditions etc., please send us the relevant information.

This includes the following information:

- Condition of the soil
- Air humidity
- Water quality
- Existing water reservoirs (lake, river, etc.)
- Rain periods
- Last harvest quantity and quality
- Other problems



PREPARING THE SOIL

Spray the soil once at the start of planting.



MIX:

- Use **1 tablet** mixed with **20-liter water** = **3000 m²**.
- Use **40 tablet** mixed with **600-800 liter** water = **1 Hectare**.
- Use **1 spoon powder (0.125g)** mixed with **20 liter** water = **3000 m²**.
- Use **40 spoon powder (5.0g)** mixed with **600-800 liter** water = **1 Hectare**.



Method: Use a Knapsack sprayer, Tractor with spray tank or Drone and apply with a mist spray.

Contact

Mobile: [+31 652 642 332](tel:+31652642332)

Email: info@greenbullone.com

Adresse

Hollandselaan 2
1213 AN Hilversum
The Netherlands



RICE & GRAIN

Begin by soaking the seed for 1 night. Soaking the seed will make the seed sprout quickly and evenly.



MIX:

- Use **1 tablet** mixed with **20-liter water** = 3000 m^2 .
- Use **40 tablet** mixed with **600-800 liter** water = 1 Hectare.
- Use **1 spoon powder (0.125g)** mixed with **20 liter** water = 3000 m^2 .
- Use **40 spoon powder (5.0g)** mixed with **600-800 liter** water = 1 Hectare.

!!! After soaking, the rice seeds should be left to dry before planting.



Method: Use a Knapsack sprayer, Tractor with spray tank or Drone and apply with a mist spray.



Spraying Schedule:

- 1st Spray: When seedlings are aged **20 - 30 days**.
- 2nd Spray: When seedlings begin to form clumps at **45 - 50 days**.
- 3rd Spray: When the rice begins to husk at **60 - 75 days**.
- 4th Spray: When rice husks mature at approximately **90 days**.

Spraying should be avoided when rice is pollinating (Flowering period), because spraying will disturb the pollination process. Efficiency Booster contributes to good root formation and stronger, resulting in strong stems, upright leaves, long pods and good weight. It can also reduce the cost of soil fertilizer by 30-50 %.



Contact

Mobile: [+31 652 642 332](tel:+31652642332)

Email: info@greenbullone.com

Adresse

Hollandselaan 2
1213 AN Hilversum
The Netherlands

RUBBER TREE



Seedling - 1 year:



MIX:

- Use **1 tablet** mixed with **20-liter water** = **3000 m²**.
- Use **40 tablet** mixed with **600-800 liter** water = **1 Hectare**.
- Use **1 spoon powder (0.125g)** mixed with **20 liter** water = **3000 m²**.
- Use **40 spoon powder (5.0g)** mixed with **600-800 liter** water = **1 Hectare**.



Method: Use a 1 Knapsack sprayer and apply with a mist.



Spraying Schedule:

Spray every 10 – 15 days or at longer intervals if the Seedling are overgrown.

- **Large Tree – 1 year & up:** Spray every **10 – 15 days** reach the fresh leaf or stem cells.
- **Small Tree:** Use the same method and spray evenly **10 – 15 days**.
- **Tall Tree:** Use the same method for spraying the trunks/bases of the tree; the rubber tree can be sprayed **10 – 15 days**.



PALM TREE

Seedling – 1 year:



MIX:

- Use **1 tablet** mixed with **20-liter water** = **3000 m²**.
- Use **40 tablet** mixed with **600-800 liter** water = **1 Hectare**.
- Use **1 spoon powder (0.125g)** mixed with **20 liter** water = **3000 m²**.
- Use **40 spoon powder (5.0g)** mixed with **600-800 liter** water = **1 Hectare**.



Method: Use a Knapsack sprayer.



Spraying Schedule:

Spray lightly and evenly every **10 - 15 days** at longer intervals if you are afraid of the seedlings becoming overgrown.

Contact

Mobile: [+31 652 642 332](tel:+31652642332)

Email: info@greenbullone.com

Adresse

Hollandselaan 2
1213 AN Hilversum
The Netherlands



Palm Tree – 1 year & up: Use the same method and spray every **10 – 15 days**; spraying the fresh cells of leaves or stems on the trunk of the palm tree would be great.

- **Small Trees:** Use the same mix and method and spray evenly.
- **Tall Trees:** Use the same ratio and method for spraying the trunks/bases of the trees



FIELD CROPS : SUGAR CANE, VAMS, CORN

Begin by soaking the seed for 30 minutes possibly along with soaking in germicides used by other farmers. Soaking the seed will make the seed sprout quickly and evenly.



MIX:

- Use **1 tablet** mixed with **20-liter water** = **3000 m²**.
- Use **40 tablet** mixed with **600-800 liter** water = **1 Hectare**.
- Use **1 spoon powder (0.125g)** mixed with **20 liter** water = **3000 m²**.
- Use **40 spoon powder (5.0g)** mixed with **600-800 liter** water = **1 Hectare**.
-



Method: Use a Knapsack sprayer, Tractor with spray tank or Drone and apply with a mist spray.



Spraying Schedule:

After Planting: **Begin spraying 1 week after planting** along with spraying other substances used by farmers spray every **15 – 20 days**.



LARGE TREES AND FRUITS

Seedling - 1 year:

Contact

Mobile: [+31 652 642 332](tel:+31652642332)

Email: info@greenbullone.com

Adresse

Hollandselaan 2

1213 AN Hilversum

The Netherlands



MIX:

- Use **1 tablet** mixed with **20-liter water** = 3000 m^2 .
- Use **40 tablet** mixed with **600-800 liter** water = 1 Hectare.
- Use **1 spoon powder (0.125g)** mixed with **20 liter** water = 3000 m^2 .
- Use **40 spoon powder (5.0g)** mixed with **600-800 liter** water = 1 Hectare.
-



Method: Knapsack sprayer and spray lightly.



Spraying Schedule:

- Large Tree 1 year & up: Use the same mix and method and spray every 10 – 15 days reach the fresh leaf or stem cells.
- Small Trees: Use the same mix and method and spray evenly.
- Tall Trees: Use the same mix and spray the same ratio and method for spraying the trunks/bases of the trees.



LEAFY GREEN VEGETABLES AND KITCHEN GARDENS

Chili, Eggplants, Luffa Gourd, Long Beans, Soybean and Cucumbers, etc.



MIX:

- Use **1 tablet** mixed with **40 liter water** = 3000 m^2 .
- Use **40 tablet** mixed with **1200-1600 liters water** = 1 Hectare.
- Use **1 spoon powder** mixed with **40 liter water** = 3000 m^2 .
- Use **40 spoon powder** mixed with **1200-1600 liters water** = 1 Hectare.



Method: Use a Knapsack sprayer, Tractor with spray tank or Drone and apply with a mist spray.



Spraying Schedule:

Spray evenly every **7 – 10 days or 1 -2 Days** before and after harvesting by observing the thickness of the plant leaves, soil density or spray whenever convenient.



FLOWERING PLANTS, ORNAMENTALS AND SEEDLINGS



MIX:

- Use **1 tablet** mixed with **40 liter water** = 3000 m^2 .

Contact

Mobile: [+31 652 642 332](tel:+31652642332)

Email: info@greenbullone.com

Adresse

Hollandselaan 2

1213 AN Hilversum

The Netherlands



- Use **40 tablet** mixed with **1200-1600 liters water** = 1 Hectare.
- Use **1 spoon powder** mixed with **40 liter water** = 3000 m².
- Use **40 spoon powder** mixed with **1200-1600 liters water** = 1 Hectare.



Method: Use a Knapsack sprayer or Drone and apply with a mist spray.



Spraying Schedule:

Spray evenly every **7 – 10 days or 1 -2 Days** after harvesting to nourish or restore the plants.



MUSHROOM (ALL TYPES)



MIX: Use 1 tablet mixed with 80 liters of water.

- Use **1 tablet** mixed with **80 liter water**
- Use **1 spoon powder** mixed with **80 liter water**

Begin with the mixing of the mushroom spawn.



Method: Knapsack sprayer



Spraying Schedule:

Spray every **5 – 7 days** after the mushrooms have started blooming

Contact

Mobile: [+31 652 642 332](tel:+31652642332)

Email: info@greenbullone.com

Adresse

Hollandselaan 2
1213 AN Hilversum
The Netherlands